

## 2020 CCS Self-Assessment Items

### ICD-10-CM DX (1-10)

1. The physician documents a severe combined immunodeficiency (SCID) due to adenosine deficiency. What diagnosis code is assigned?
  - a. D81.30
  - b. D81.31
  - c. D81.32
  - d. D81.39
  
2. A patient is admitted for a gastrojejunostomy tube change due to a clogged gastrojejunostomy (GJ) tube. What diagnosis code is assigned for the GJ tube change?
  - a. K94.23
  - b. K94.29
  - c. K94.20
  - d. K94.39
  
3. A 90-year-old man has a history of COPD and emphysema. He presents to the Emergency Department with COPD exacerbation. What diagnosis code(s) should be assigned?
  - a. J43.9
  - b. J44.9, J44.1
  - c. J44.1
  - d. J44.0
  
4. The patient presents for with nosocomial gram-negative bacilli sepsis/blood stream infection, due to a midline catheter. What diagnosis codes are assigned?
  - a. T82.7XXA, T80.211A
  - b. T82.7XXA, T80.219A
  - c. T82.7XXA, A41.89
  - d. T82.7XXA, A41.59
  
5. The provider documented a diagnosis of choledochoduodenal fistula with choledocholithiasis, cholecystitis and obstruction. What diagnosis codes are assigned?
  - a. K80.40, K83.1
  - b. K80.41, K83.3
  - c. K80.43, K83.1
  - d. K80.45, K83.3
  
6. A patient is admitted with a midline catheter developed cellulitis of the right upper arm secondary to the midline catheter. What diagnosis codes should be assigned?
  - a. T82.7XXA, L03.114
  - b. T82.7XXD, L03.113
  - c. T82.7XXA, L03.113
  - d. T82.7XXA, L03.115
  
7. A 65-year-old man presents to the Emergency Department (ED) with an elevated troponin level. An acute myocardial infarction is ruled-out. What diagnosis code should be assigned?
  - a. R79.9
  - b. R74.9

- c. R77.9
- d. R79.89

8. A 23-year-old patient presents to the Emergency Department (ED) with an acute on chronic complex left medial meniscus tear/injury? What diagnosis code should be assigned?

- a. S83.232A
- b. S83.232D
- c. S83.242A
- d. S83.242D

9. A patient is admitted with a neuroendocrine tumor of the colon with metastatic disease to the liver. What diagnosis codes should be assigned?

- a. C7A.8, C7B.00
- b. C7A.098, C7B.02
- c. C7A.098, C7B.09
- d. C7A.8, C7B.8

10. A 45-year-old female with emphysema presents to the Emergency Department with melena and hematemesis. She was diagnosed with acute on chronic blood loss anemia due to a bleeding duodenal ulcer. What is the diagnosis code assignment for the acute on chronic blood loss anemia?

- a. D50.0
- b. D62
- c. D50.8
- d. D50.9

## ICD-10-PCS (11-17)

11. A patient with lumbar spinal stenosis and spondylolisthesis with scoliosis has a decompressive laminectomy at L2, L3 and L4 followed by spinal fusion of L3-4 and L4-5 with interbody cages. What is/are the correct root operation(s)?

- a. Code only Fusion for the interbody cages
- b. Code only Insertion for the interbody cages
- c. Code Excision for the decompressive laminectomy and Fusion for the interbody cages
- d. Code Release for the decompressive laminectomy and Fusion for the interbody cages

12. A patient with carcinoma of the pancreas presents for excision. The abdomen is prepped and draped. Incision is made in the upper abdomen and the central portion of the pancreas is excised. The pancreatic neck and anterior body are then anastomosed to the posterior wall of the stomach. The anastomosis is reinforced with an omental flap. What is/are the correct ICD-10-PCS code(s) for this procedure?

- a. 0FBG0ZZ
- b. 0FBG0ZZ, 0DU607Z
- c. 0FBG0ZZ, 0F1G0Z3
- d. 0FBG0ZZ, 0F1G0Z3, 0DU607Z

13. During an upper endoscopy, a large chronic perforation was found on the posterior wall of the stomach. A wound vacuum assisted closure sponge was placed via endoscope onto the perforation and liquid was suctioned into the wound VAC. What is/are the correct ICD-10-PCS code(s) for this procedure?

- a. 0DQ68ZZ
- b. 0DQ68ZZ, 0D968ZZ
- c. 0DH68YZ
- d. 0DU68JZ

14. Patient with a pathologic compression fracture of L4 presents for Kiva® kyphoplasty. A needle was advanced across the pedicle of L4 and into the vertebral body. The Kiva® implant was deployed and bone cement injected into the implant to reinforce the fractured vertebral body. What is the correct ICD-10-PCS code for this procedure?
- a. 3E0V3GC
  - b. 0QU03JZ
  - c. 0QS034Z
  - d. 0SS034Z
15. Patient with a pilonidal cyst and sinus with multiple recurrent abscesses presents for treatment. The resectoscope was placed in cautery mode and the pilonidal cyst and sinus were cut out using cautery. The abscess cavity was cored out and the nonhealing sinus tract removed. The resectoscope was used again in cautery mode to remove the chronic inflammatory tissue and simultaneously cauterize any hair-bearing area to prevent recurrence. What is the correct ICD-10-PCS code for this procedure?
- a. 0JB93ZZ
  - b. 0JC93ZZ
  - c. 0JD93ZZ
  - d. 0J593ZZ
16. Patient with chronic COPD presents for placement of endobronchial one-way valves into the left lower lobe bronchus via bronchoscope. What is the correct ICD-10-PCS code for this procedure?
- a. 0BUB8JZ
  - b. 0BHB8GZ
  - c. 0BVB8DZ
  - d. 0BLB8DZ
17. Patient with peritoneal metastasis presents for hyperthermic intraperitoneal chemotherapy with mitomycin. Inflow and outflow cannulas were placed in the patient's peritoneal cavity and mitomycin was perfused for two hours. What is the correct ICD-10-PCS code for this procedure?
- a. DWY38ZZ
  - b. 3E0M30Y
  - c. DWY58ZZ
  - d. 3E0M3HZ

### CPT 10 CPT Questions (18-27)

18. An existing percutaneous gastrostomy tube requires removal and replacement of the tube due to a clogged device. What is the correct CPT code?
- a. 43761
  - b. 43762
  - c. 43763
  - d. 43773
19. A 63-year-old female undergoes bladder neck suspension. The surgeon uses tension-free vaginal tape for bladder neck support. What is/are the correct CPT code(s)?
- a. 57260
  - b. 57288
  - c. 57288, 52000-59
  - d. 57289
20. An esophageal sphincter augmentation device was placed during a laparoscopic hiatal hernia repair. What is/are the correct CPT code(s)?

- a. 43280
  - b. 43281
  - c. 43280, 43284-59
  - d. 43284
21. A 24-year-old patient has bronchodilation responsiveness testing performed. What is/are the correct CPT code(s)?
- a. 94010, 99070
  - b. 94060, 99070
  - c. 94060
  - d. 94070
22. A PICC lined is placed with imaging guidance for a 4-year-old requiring long-term IV antibiotic therapy. What is the correct CPT code?
- a. 36568
  - b. 36572
  - c. 36573
  - d. 36570
23. A 34-year-old female presents for a bilateral breast MRI. She is status post right mastectomy and due to genetic mutation is at high risk for breast cancer. MRI is with and without contrast and includes the use of computer-aided detection. What is the correct CPT code?
- a. 77046
  - b. 77047
  - c. 77048
  - d. 77049
24. Patient underwent an ORIF for a fracture of the second metatarsal head fracture. What is the correct CPT code?
- a. 28456
  - b. 28465
  - c. 28475
  - d. 28485
25. Patient undergoes placement of a Stimware stimulator electrode at the level of the L4 vertebral body through the epidural space. What is/are the correct CPT code(s)?
- a. 63650, C1778
  - b. 63655, C1778
  - c. 63685, C1778
  - d. 95970
26. Patient underwent removal of inferior scleral sutures by the original surgeon as a follow-up to a repair of a ruptured globe. The procedure was performed under MAC anesthesia. What is the correct CPT code?
- a. 15850
  - b. 15851
  - c. 17999
  - d. 66999
27. Patient underwent bilateral laser reduction of a vocal cord mass via direct microlaryngoscopy. What is the correct CPT code?
- a. 31535

- b. 31536
- c. 31540
- d. 31541

### General 3 Questions

28. In ICD-10-CM, Body Mass Index age range for adults changed from 21 years of age or older to:
- a. 17 years of age or older
  - b. 18 years of age or older
  - c. 19 years of age or older
  - d. 20 years of age or older
29. The Medicare Code Editor Age conflict edit for Maternity Diagnoses was revised from an age range of 12 to 55 years to 9 to 64 years age.
- a. True
  - b. False
30. When assigning a code for physeal fractures which of the following is the appropriate convention?
- a. Assign both the type of physeal fracture and a code to identify the specific bone that is fractured
  - b. Assign only the code identifying the type of physeal fracture.
  - c. Assign only the code identifying the specific bone that is fractured.
  - d. Assign all physeal fractures as Type III